

3. [Maximum mark: 5]

Let $f'(x) = \frac{8x}{\sqrt{2x^2 + 1}}$. Given that $f(0) = 5$, find $f(x)$.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....


.....

.....




4. [Maximum mark: 5]

The following diagram shows the graph of $y = f(x)$. The graph has a horizontal asymptote at $y = -1$. The graph crosses the x -axis at $x = -1$ and $x = 1$, and the y -axis at $y = 2$.



On the following set of axes, sketch the graph of $y = [f(x)]^2 + 1$, clearly showing any asymptotes with their equations and the coordinates of any local maxima or minima.



6. [Maximum mark: 8]

(a) Show that $\log_9(\cos 2x + 2) = \log_3 \sqrt{\cos 2x + 2}$. [3]

(b) Hence or otherwise solve $\log_3(2 \sin x) = \log_9(\cos 2x + 2)$ for $0 < x < \frac{\pi}{2}$. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....



Turn over

9. [Maximum mark: 8]

The function f is defined by $f(x) = e^{2x} - 6e^x + 5$, $x \in \mathbb{R}$, $x \leq a$. The graph of $y = f(x)$ is shown in the following diagram.



- (a) Find the largest value of a such that f has an inverse function. [3]
- (b) For this value of a , find an expression for $f^{-1}(x)$, stating its domain. [5]

(This question continues on the following page)



Do **not** write solutions on this page.

Section B

Answer **all** questions in the answer booklet provided. Please start each question on a new page.

10. [Maximum mark: 16]

Let $f(x) = \frac{\ln 5x}{kx}$ where $x > 0, k \in \mathbb{R}^+$.

(a) Show that $f'(x) = \frac{1 - \ln 5x}{kx^2}$. [3]


The graph of f has exactly one maximum point P.

(b) Find the x -coordinate of P. [3]

The second derivative of f is given by $f''(x) = \frac{2 \ln 5x - 3}{kx^3}$. The graph of f has exactly one point of inflexion Q.

(c) Show that the x -coordinate of Q is $\frac{1}{5}e^{\frac{3}{2}}$. [3]

The region R is enclosed by the graph of f , the x -axis, and the vertical lines through the maximum point P and the point of inflexion Q.



(d) Given that the area of R is 3, find the value of k . [7]



Do **not** write solutions on this page.

11. [Maximum mark: 18]

- (a) Express $-3 + \sqrt{3}i$ in the form $re^{i\theta}$, where $r > 0$ and $-\pi < \theta \leq \pi$. [5]

Let the roots of the equation $z^3 = -3 + \sqrt{3}i$ be u, v and w .

- (b) Find u, v and w expressing your answers in the form $re^{i\theta}$, where $r > 0$ and $-\pi < \theta \leq \pi$. [5]

On an Argand diagram, u, v and w are represented by the points U, V and W respectively.

- (c) Find the area of triangle UVW . [4]

- (d) By considering the sum of the roots u, v and w , show that $\cos \frac{5\pi}{18} + \cos \frac{7\pi}{18} + \cos \frac{17\pi}{18} = 0$. [4]

12. [Maximum mark: 21]

The function f is defined by $f(x) = e^{\sin x}$.

- (a) Find the first two derivatives of $f(x)$ and hence find the Maclaurin series for $f(x)$ up to and including the x^2 term. [8]

- (b) Show that the coefficient of x^3 in the Maclaurin series for $f(x)$ is zero. [4]

- (c) Using the Maclaurin series for $\arctan x$ and $e^{3x} - 1$, find the Maclaurin series for $\arctan(e^{3x} - 1)$ up to and including the x^3 term. [6]

- (d) Hence, or otherwise, find $\lim_{x \rightarrow 0} \frac{f(x) - 1}{\arctan(e^{3x} - 1)}$. [3]

